

Patterns and rules surround us.

For instance, the dry and wet seasons are alternating throughout the entire year; the stripes in the petals of a flower are arranged systematically; funds in a regular savings account earn interest monthly, and even your daily morning routine follows a well-defined pattern.

Quantities and numbers also follow rules and patterns. For instance, counting by 2s is one of the ways to define a pattern for numbers.

In this review, we're going to focus on **sequence** - a set of numbers following a certain pattern. Once we've unraveled its important concepts, we will deal with **series** or the sum of these patterned numbers.

## What is a Sequence?

Look at this given set of numbers:

5, 10, 15, 20, 25, 30, \_\_\_\_\_.

*What number must be placed in the blank?*

Yes, it should be 35.

It's pretty easy to guess the missing number. By observing the numbers above, you can deduce that the next number can be obtained by adding 5 to 30. I'm sure you're able to do this because you know that the numbers follow a particular pattern (i.e., adding 5 to the previous number to get the next one).

The set of numbers above (5, 10, 15, 20, 25, 30, 35) is an example of a **sequence**.

A sequence is a set of numbers that follow a specific pattern. As we have mentioned earlier, the pattern of this sequence is that it adds 5 to the previous number to get the succeeding one ( $5 + 5 = 10$ ;  $10 + 5 = 15$ ;  $15 + 5 = 20 \dots$ )

**Sample Problem:** What is the missing number in 72, 63, 54, 45, \_\_\_\_\_?

**Solution:** The sequence above follows a specific pattern. It subtracts 9 from the previous number to obtain the succeeding one ( $72 - 9 = 63$ ;  $63 - 9 = 54 \dots$ ). Therefore, to get the missing number:  $45 - 9 = 36$ .

Hence, the answer is 36.

### Terms of a Sequence

Each element in a sequence is called a term. Thus, in the sequence 5, 10, 15, 20, 25, 30, 35, each number is a term of the sequence. Moreover, 5 is called the **first term**, 10 is the **second term**, 15 is the **third term**, and so on.

### Notations in Sequence

Generally, the first term of a sequence can be expressed as  $a_1$ . The subscript "1" indicates that it is the first term of a particular sequence.

For example, in the sequence 5, 10, 15, 20, 25, 30, 35, the first term is 5. Therefore,  $a_1 = 5$ .

Now, if the first term of a sequence is expressed as  $a_1$ , then the second term must be  $a_2$ , the third term must be  $a_3$ , the fourth term must be  $a_4$ , and so on. Thus, the  $n$ th term of a sequence is written as  $a_n$ .

Hence, in general, a sequence can be written as:

$$a_1, a_2, a_3, a_4 \dots, a_n$$


**Sample Problem:** Determine  $a_2$  and  $a_3$  of 2, 5, 8, 11, 14, 17.

**Solution:**  $a_2$  and  $a_3$  refer to the second and third terms of the given sequence, respectively. Since the second term of the sequence is 5 while the third term is 8, then  $a_2 = 5$  and  $a_3 = 8$ .

### Rule of a Sequence

Let's go back to our first example: 5, 10, 15, 20, 25, 30, 35,...

We have mentioned earlier that the pattern or rule of this sequence is that it adds 5 to a previous term to obtain the next one:

A light blue rectangular box containing a sequence of numbers and equations. The numbers 5, 10, 15, 20 are in black, and the numbers 5, 5, 5, 5 are in red. The equations are 5 + 5 = 10, 10 + 5 = 15, 15 + 5 = 20, and 20 + 5 = 25. The FilipiKnow logo is in the top right corner of the box.


Using a formula or an equation, we can provide a more precise and concise description of the rule of the sequence.

For instance, the rule of the sequence 5, 10, 15, 20, 25, 30, 35 can be expressed as  $a_n = 5n$  where  $a_n$  is the  $n$ th term of the sequence. Note that the  $n$ th term means any term of the given sequence.

If we let  $n = 1$ , then by substitution:  $a_{(1)} = 5(1) = 5$ . This means that the first term of the sequence defined by the rule  $a_n = 5n$  is 5.

If  $n = 2$ , then  $a_{(2)} = 5(2) = 10$ . This means that the second term of the sequence defined by the rule  $a_n = 5n$  is 10.

Similarly, if  $n = 5$ , then  $a_{(5)} = 5(5) = 25$ . This means that the fifth term of the sequence defined by the rule  $a_n = 5n$  is 25.


$$a_n = 5n$$

**If  $n = 1$ :**

$$a_n = 5n \rightarrow a_1 = 5(1) = 5$$

**If  $n = 2$ :**

$$a_n = 5n \rightarrow a_2 = 5(2) = 10$$

**If  $n = 5$ :**

$$a_n = 5n \rightarrow a_5 = 5(5) = 25$$

**Sample Problem 1:** The sequence 5, 10, 15, 20, 25, 30, 25 is defined by the equation  $a_n = 5n$ . What is the 100th term of the sequence?

**Solution:** To find the 100th term of the sequence, which is defined by the equation  $a_n = 5n$ , we let  $n = 100$ :

$$a_{100} = 5(100) = 500$$

Therefore, the 100th term of the sequence is 500.

**Sample Problem 2:** A sequence is defined by the rule  $a_n = 7n - 1$ . What is the first term of this sequence? How about its 20th term?

**Solution:** To find the first term of the sequence defined by the rule  $a_n = 7n - 1$ , we let  $n = 1$ :

$$a_1 = 7(1) - 1 = 6 - 1 = 5$$

Therefore, the first term of the sequence is 5.

On the other hand, to find the 20th term, we let  $n = 20$ :

$$a_{20} = 7(20) - 1 = 140 - 1 = 139$$

Thus, the 20th term of the sequence is 139.

**Sample Problem 3:** Given the sequence with a rule defined by the equation  $a_n = 5 - 2n$ , what is the 15th term?

**Solution:** To find the 15th term, we let  $n = 15$ :

$$a_{15} = 5 - 2(15) = 5 - 30 = -25$$


Thus, the 15th term of the sequence is -25.

## Types of Sequence

### 1. Arithmetic Sequence

An arithmetic sequence is a sequence wherein every succeeding term is obtained by adding a **fixed** number to the previous term.

Our previous example (i.e., 5, 10, 15, 20, 25, 30, 35) is an example of an arithmetic sequence because its succeeding term is derived by adding a fixed number (which is 5) to the previous term.



The fixed number added to every term to get the succeeding one is called the **common difference**.


For instance, the sequence 5, 10, 15, 20, 25, 30, and 35 has a common difference of 5.

**Sample Problem 1:** Which of the following is/are arithmetic sequence(s)?

1. 8, 10, 12, 14, 16, 18, ...
2. 15, 12, 9, 6, 3, 0, ...
3. 2, 4, 8, 16, 32, ...


**Solution:**

1. Every succeeding term of this sequence is obtained by adding 2 to the previous term.




Therefore, this sequence is an arithmetic sequence with a common difference of 2.

2. This sequence works this way: it subtracts 3 from the previous term to obtain the next one. Note that subtracting 3 from a number also means adding -3 to a number.



Since we are still adding a fixed number (-3) to every term in the sequence, this is an arithmetic sequence. Thus, this is an example of an arithmetic sequence with a common difference of -3.

- This sequence is not an arithmetic sequence because the number added to each term to obtain the next one is not fixed or constant.



Therefore, only sequences in A and B are arithmetic.

**Sample Problem 2:** Determine the common difference of the following arithmetic sequence.

1. 11, 22, 33, 44, 55, ...
2. -8, -5, -2, 1, 4, ...
3.  $\frac{2}{3}$ , 1,  $\frac{4}{3}$ ,  $\frac{5}{3}$ , ...

**Solution:** To find the common difference ( $d$ ) of an arithmetic sequence, we just pick any term in the given sequence and subtract from it the term that precedes it. Mathematically,

$$d = a_n - a_{n-1}$$

Where  $d$  is a common difference,  $a_n$  is any term of the sequence ( $n$ th term), and  $a_{n-1}$  is the term that precedes  $a_n$ .

1. To identify the common difference in this arithmetic sequence, we can subtract from any of its terms the term that precedes it

Let's try 33; the term preceding it is 22. Hence, the common difference is:

$$d = 33 - 22 = 11$$

We can still get a common difference if we try another term. Let's try 44; the term preceding it is 33:

$$d = 44 - 33 = 11$$

Hence, the common difference is 11.

2. Using the same method we used in the previous item, we can pick any term and subtract from it the term preceding it:

$$d = -2 - (-5) = -2 + 5 = 3$$

Hence, the common difference is 3.

3. To find the common difference, we just pick any term and subtract from it the term that precedes it:



$$d = 5/3 - 4/3 = 1/3$$


Therefore, the common difference is  $1/3$ .

### Finding the $n$ th Term of an Arithmetic Sequence

Suppose we have this arithmetic sequence: 6, 9, 12, 15, 18, ... and we want to find its 25th term. What will you do to find this term?

If you're planning to add 3 to each previous term until you reach the 25th term, then you'll waste a lot of your precious time. Perhaps you're good at arithmetic, and you can reach the 25th term in a couple of seconds. However, what if you're looking for the 100th, 250th, or 500th term? Enumerating the terms one by one becomes inconvenient and time-consuming in this case.

Luckily, mathematicians have done the heavy work and given us a general formula that we can use to find any term of an arithmetic sequence without listing each term manually. This formula is presented in the image below:

A light blue rectangular box containing the formula  $a_n = a_1 + (n - 1)d$  in bold black text. A small FilipiKnow logo is in the top right corner of the box.
$$a_n = a_1 + (n - 1)d$$

Here is the meaning of the variables in the formula above

- $a_n$  is any term of the sequence; this is the term you're looking for
- $a_1$  is the first term of the given sequence
- $n$  is the number of the term you're looking for (e.g., if you're looking for the 25th term, then  $n = 25$ ). If the last term of an arithmetic sequence is specified,  $n$  also represents the total number of terms in the sequence.
- $d$  is the common difference of the arithmetic sequence

We will not discuss how this formula is derived because it's quite long and complex.

For now, let's answer some examples to understand better how to use this formula.

**Sample Problem 1:** Determine the 25th term of the sequence 6, 9, 12, 15, 18, ...

**Solution:** The formula to find any term of an arithmetic sequence is defined as:

$$a_n = a_1 + (n - 1)d$$

- The first term of the sequence is 6, so we have  $a_1 = 6$
- Since we are looking for the 25th term, then we have  $n = 25$
- The common difference ( $d$ ) of the given sequence is 3 (e.g.,  $9 - 6 = 3$ )

Substituting the values we have derived:

$$a_n = a_1 + (n - 1)d$$

$$a_{25} = 6 + (25 - 1)3$$

$$a_{25} = 6 + (24)3$$

$$a_{25} = 6 + 72$$

$$a_{25} = 78$$

Thus, the 25th term of the sequence is 78.

Let us verify if the formula works by listing each term of the given sequence manually until the 25th term:

6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57, 60, 63, 66, 69, 72, 75, **78**

The list above shows that indeed, 78 is the 25th term of the given sequence.

**Sample Problem 2:** Identify the 54th term of the sequence 17, 28, 39, 40, 51, 62, ...

**Solution:** The formula to find any term of an arithmetic sequence is defined as:

$$a_n = a_1 + (n - 1)d$$

- The first term of the sequence is 17, so we have  $a_1 = 17$
- Since we are looking for the 54th term, then we have  $n = 54$
- The common difference (d) of the given sequence is 11 ( $28 - 17 = 11$ )

Substituting the values we have derived:

$$a_{54} = a_1 + (n - 1)d$$

$$a_{54} = 17 + (54 - 1)11$$

$$a_{54} = 17 + (53)11$$

$$a_{54} = 17 + 583$$

$$a_{54} = 600$$

Thus, the 54th term of the sequence is 600.

**Sample Problem 3:** Selena plans to save money for 30 days. On the first day, she saved PHP 20. On the second day, she saved PHP 22. On the third day, she saved PHP 24. On the fourth day, she saved PHP 26. How much will Selena save on the 30th day?

**Solution:** The manner in which Selena saves money follows an arithmetic sequence of

20, 22, 24, 26, ... This sequence has a common difference of 2 ( $d = 22 - 20 = 2$ ).

To find how much Selena will save on the 30th day, we need to find the 30th term of the given sequence. The formula to find any term of an arithmetic sequence is defined as:

$$a_n = a_1 + (n - 1)d$$

- The first term of the sequence is 20, so we have  $a_1 = 20$
- Since we are looking for the 30th term, we have  $n = 30$
- The common difference (d) of the given sequence is 2

Substituting the values we have derived:

$$a_{30} = a_1 + (n - 1)d$$

$$a_{30} = 20 + (30 - 1)2$$

$$a_{30} = 20 + (29)2$$

$$a_{30} = 20 + 58$$

$$a_{30} = 78$$

Thus, the 30th term of the sequence is 78; Selena will save PHP 78 on the 30th day.

**Sample Problem 4:** How many numbers from 1 to 100 are divisible by 4?


**Solution:** At first glance, the given problem seems unrelated to the concept of the arithmetic sequence. However, this problem is solvable by applying what we have learned about this type of sequence.

If a number is divisible by 4, it is also a multiple of 4. For instance, 16 is a multiple of 4, so it also means that it is also divisible by 4.

To identify the numbers that are multiple (and divisible) by 4 between 1 to 100, we just count by 4s:

4, 8, 12, 16, 20, 24, ...

Note that the list of numbers above is an arithmetic sequence with a common difference of 4.



Thus, to find how many numbers from 1 - 100 are divisible by 4, we need to identify how many terms there are in the sequence.

However, we cannot identify the total number of terms contained in the sequence without figuring out its last term. Note that the last number between 1 to 100 that is divisible by 4 is 96. This also means that the last term of our sequence should be 96.

Hence, the arithmetic sequence that corresponds to all numbers between 1 to 100 divisible by 4 is 4, 8, 12, 16, 20, 24, ..., 96. We can now identify how many terms there are in this sequence:

The formula for identifying any term of a given arithmetic sequence is

$$a_n = a_1 + (n - 1)d$$

However, this time we are not looking for  $a_n$  but  $n$ . Note that if the last term of a sequence is specified, then  $n$  may represent the total number of terms in a sequence. Since the sequence 4, 8, 12, 16, 20, 24, ..., 96 ends with 96, we have a sequence with a specified last term. So,  $n$  in this problem serves as the total number of terms in the sequence (the total number of numbers from 1 - 100 that are divisible by 4).

- Since 4 is the first number in the sequence, then we have  $a_1 = 4$
- Since we are interested in finding the number of terms in 4, 8, 12, 16, 20, ..., 96 or the value of " $n$ ," the value of  $a_n$  should be the last term which is 96.
- The common difference ( $d$ ) is 4

Substituting these values in the formula, we can now solve for  $n$ :

$$a_n = a_1 + (n - 1)d$$

$$96 = 4 + (n - 1)4$$

Let us now do some algebra to solve the value of  $n$ :

$$96 = 4 + (n - 1)4$$

$$96 = 4 + 4n - 4 \quad \text{Distributive Property}$$

$$96 = 4n \quad \text{Combining like terms}$$

$$96/4 = 4n/4 \quad \text{Dividing both sides of the equation by 4}$$

$$24 = n$$

$$n = 24$$

Since  $n$  stands for the number of terms in the sequence 4, 8, 12, 16, 20, 24, ... 96, there are 24 terms in this sequence. This also implies that there are 24 numbers between 1 to 100 that are divisible by 4.

**Sample Problem 5:** The first term of an arithmetic sequence is -9 while its 12th term is 1. What is the common difference between this arithmetic sequence?

**Solution:** We have  $a_1 = -9$  since -9 is the first term of the sequence. On the other hand,  $a_{12}$  (the  $a_n$  in this case) is 1. Furthermore, we have  $n = 12$ . Our goal is to solve for the common difference or  $d$ .

$$a_n = a_1 + (n - 1)d$$

$$1 = -9 + (12 - 1)d \quad \text{By substitution}$$

$$1 = -9 + 11d$$

$$1 + 9 = 11d \quad \text{Transposition method}$$

$$10 = 11d$$

$$10/11 = 11d/11 \quad \text{Dividing both sides of the equation by 11}$$

$$d = 10/11$$


Thus, the common difference in the sequence in this problem is  $10/11$ .

## 2. Geometric Sequence

In an arithmetic sequence, there's a fixed number added to each term to obtain the succeeding term (called the common difference). Meanwhile, **a geometric sequence has each term multiplied by a fixed number to obtain the next one.**

Look at this sequence: 3, 6, 12, 24, 48, ...

Unlike in an arithmetic sequence, there is no common difference. Instead, each term is multiplied by 2 to obtain the next one.



Since this sequence is characterized by multiplying a term by 2 to get the next one, this is an example of a geometric sequence.

In a geometric sequence, each term is multiplied by a fixed number to get the next term; the fixed number is called the **common ratio** (denoted by the letter  $r$ ).

This means that the common ratio of the sequence 3, 6, 12, 24, 48, ... is 2.

**Sample Problem 1:** Which of the following is/are geometric sequence(s)?

1. 1, 4, 7, 10, 13, ...
2. 2, 6, 18, 54, ...
3. 9, 11, 14, 18, ...

**Solution:**

1. This sequence has a common difference of 3. Thus, it is not a geometric sequence.
2. This sequence has a common ratio of 3. This means that to obtain the succeeding term, we multiply the previous term by 3 (e.g.,  $2 \times 3 = 6$ ;  $6 \times 3 = 18$ ). Hence, this is a geometric sequence.
3. This sequence is neither arithmetic nor geometric sequence. It has no common difference or common ratio.

**Sample Problem 2:** Identify the common ratio of the following sequence.

1. 1, 5, 25, 125, 625, ...
2. 3, -6, 12, -24, 48, ...

**Solution:**

Here's a simple trick to identify the common ratio of a geometric sequence. Pick any term from the sequence then divide it by the term that precedes it. Let us apply this trick to find the common ratio of the sequence:

1. Let us pick 25. The term that precedes it is 5. Hence, the common ratio is:



$$r = 25/5 = 5$$

2. Let us pick -24. The term that precedes it is 12. Thus, the common ratio is:

$$r = -24/12 = -2$$

Again, we can find the common ratio of a geometric sequence by picking any term from the sequence (except the first term), then dividing it by the term that precedes it. Mathematically,

$$r = \frac{a_n}{a_{n-1}}$$

Where  $r$  is the common ratio,  $a_n$  is the  $n$ th term (any term of the sequence), and  $a_{n-1}$  is the term that precedes  $a_n$ .

#### Finding the $n$ th Term of a Geometric Sequence

Like in an arithmetic sequence, we can also determine any term of a geometric sequence using a formula. Suppose we want to identify the 10th term of 3, 6, 12, 24, ... we can use the formula below to find it without listing the terms.

The formula  $a_n = a_1 r^{n-1}$  is displayed in a light blue rectangular box. The text "FilipiKnow" is visible in the top right corner of the box.
$$a_n = a_1 r^{n-1}$$

Where:

- $a_n$  is any term of the sequence; this is the term you're looking for
- $a_1$  is the first term of the given sequence

- **n** is the number of the term you're looking for (For example, if you're looking for the 25th term,  $n = 25$ )
- **r** is the common ratio of the geometric sequence

**Sample Problem 1:** Find the 10th term of the geometric sequence 3, 6, 12, 24, ...

**Solution:** The first term of the sequence is 3. Thus, we have  $a_1 = 3$ . Since we are looking for the 10th term or  $a_{10}$ , we have  $n = 10$ .

The common ratio of the sequence is  $r = 6/3 = 2$ .

Using the formula for the  $n$ th term of a geometric sequence:  $a_n = a_1 r^{n-1}$

$$a_n = a_1 r^{n-1}$$

$$a_{10} = 3(2)^{10-1} \quad \text{By substitution}$$

$$a_{10} = 3(2)^9$$

$$a_{10} = 3(512) \quad 2^9 = 512$$

$$a_{10} = 1536$$

Therefore, the 10th term of the geometric sequence is 1,536.

**Sample Problem 2:** What is the 8th term of the geometric sequence  $1/16, 1/8, 1/4, 1/2, \dots$

**Solution:** The first term of the sequence is  $1/16$ . Thus, we have  $a_1 = 1/16$ . Since we are looking for the 8th term or  $a_8$ , we have  $n = 8$ .

The common ratio of the sequence is

$$r = \frac{1}{8} \div \frac{1}{16} = \frac{1}{8} \times \frac{16}{1} = \frac{16}{8} = 2$$

You can [review our module about how to divide fractions](#) to learn how we're able to obtain the answer above.

Using the formula for the  $n$ th term of a geometric sequence:  $a_n = a_1 r^{n-1}$

$$a_n = a_1 r^{n-1}$$

$$a_8 = 1/16(2)^{8-1} \quad \text{By substitution}$$

$$a_8 = 1/16(2)^7$$

$$a_8 = 1/16(128) \quad 2^7 = 128$$

$$a_8 = 128/16$$

$$a_8 = 8$$

Therefore, the 8th term of the geometric sequence is 8.

### 3. Harmonic Sequence

Before we proceed to discuss what a harmonic sequence is, take a look first at this arithmetic sequence: 4, 7, 10, 13, 16, 19, ...


Let us take the reciprocal of each term of the arithmetic sequence above:

$$\frac{1}{4}, \frac{1}{7}, \frac{1}{10}, \frac{1}{13}, \frac{1}{16}, \frac{1}{19}, \dots$$

The set of fractions you see above is an example of a harmonic sequence.

A harmonic sequence is derived from the reciprocals of the terms of an arithmetic sequence. This also means that the reciprocal of the terms of a harmonic sequence form an arithmetic sequence.

Going back to our example of a harmonic sequence above, taking the reciprocal of each term forms an arithmetic sequence with a common difference of 3:



This type of sequence is called “harmonic” because it is said that ancient Greek mathematicians like Pythagoras used this particular type of numerical pattern to devise a “mathematical” theory that could explain musical harmony.

**Sample Problem:** Is  $1/9, 1/7, 1/5, 1/3, \dots$  a harmonic sequence or not?

**Solution:** If we take the reciprocal of each term of the sequence above, then:

$$9, 7, 5, 3, \dots$$


It’s clear that the sequence above is an arithmetic sequence because it has a common difference of -2. Therefore, the original sequence is a harmonic sequence.

### 4. Fibonacci Sequence

A Fibonacci sequence is a sequence wherein the succeeding term is obtained by adding two previous terms. It's named after Leonardo "Fibonacci" of Pisa, an Italian mathematician who first discussed it during the early 1200s.

Here's an example of a Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, ...

In the sequence above, any term (except for the first and second terms) is obtained by adding two previous terms. For instance, to obtain 2, we add two terms preceding it which are 1 and 1 (i.e.,  $1 + 1 = 2$ ). Similarly, to obtain 8, we add the two terms before it (i.e.,  $3 + 5 = 8$ ).



Therefore, the  $n$ th term of a Fibonacci sequence can be expressed mathematically as follows:

$$a_n = a_{n-1} + a_{n-2}$$

where  $a_n$  is any term of the sequence, with  $a_{n-1}$  and  $a_{n-2}$  corresponding to the two terms that precede  $a_n$ .

**Sample Problem:** What is the missing number in this sequence: 3, 3, 6, 9, 15, 24, \_\_\_\_?

**Solution:** The sequence above is a Fibonacci sequence because each succeeding term is obtained by adding two previous terms.

Hence, to find the missing number, we simply add 15 and 24:  $15 + 24 = 39$ .


Thus, the missing number is 39.

## What is a Series?

In mathematics, a series is a sum of all terms of a particular sequence.

For instance, consider this sequence: 3, 5, 7, 9, 11, ...

If we add all of the terms of the sequence above, then it becomes a series:  $3 + 5 + 7 + 9 + 11 + \dots$

A light blue rectangular box containing text. At the top right is a small FilipiKnow logo. The word "Sequence" is written in bold red text. Below it, the numbers "3, 5, 7, 9, 11, ..." are listed in bold black text. The word "Series" is written in bold red text. Below it, the expression "3 + 5 + 7 + 9 + 11 ..." is written in bold black text.

**Sequence**  
**3, 5, 7, 9, 11, ...**

**Series**  
**3 + 5 + 7 + 9 + 11 ...**

Hence, if  $a_1, a_2, a_3, \dots, a_n$  represents the terms of a sequence, then a series  $S$  is defined as:

$$S = a_1 + a_2 + a_3 + \dots + a_n$$

**Sample Problem:** Transform the sequence  $-4, -2, 0, 2, 4, 6, \dots$  into a series.

**Solution:** To turn the sequence above into a series, we simply express it as a sum of its terms:

$$(-4) + (-2) + 0 + 2 + 4 + 6 + \dots$$

### Partial Sum and Infinite Series

A mathematical series can either be a partial sum or an infinite sum (series).


A series is a **partial sum** if we are just adding a part of a sequence. That is, we are just adding a finite number of terms of the sequence.

For example, if we are just interested in adding the first 10 terms of the sequence  $3, 5, 7, 9, \dots$ , the series  $3 + 5 + 7 + 9 \dots + 21$  is a partial sum of the sequence (21 is the 10th term of the given sequence).

On the other hand, if we are adding a limitless number of terms in a sequence, then we're now dealing with an infinite series. For instance, adding all terms of the sequence  $3, 5, 7, 9, \dots$  up to infinity is an example of an infinite series.

An **infinite series** is a series where an infinite number of terms of a sequence are added.

You might find it ridiculous why someone would add the terms of a sequence like  $3, 5, 7, 9, \dots$  up to infinity. We cannot arrive at an exact sum because we are dealing with an infinite number of addends. The sum will just get larger and larger without reaching a definite end.


$$3 + 5 + 7 + 9 + 11 \dots$$
$$3 + 5 = 8 + 7 = 15 + 9 = 24 + 11 = 35 + \dots$$

However, whether you believe it or not, some infinite series “converge” to an exact result. That is, it is possible to obtain an exact sum by adding an endless number of terms for some sequences. For instance, if you add all terms of the sequence  $\frac{1}{2}$ ,  $\frac{1}{4}$ ,  $\frac{1}{8}$ ,  $\frac{1}{16}$ , .. up to infinity, the sum will be 1.

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots = 1$$

I know this thing seems counterintuitive and mind-boggling. Don't worry as we will shed light on this confusing thing later as we explain the type of infinite series with an “exact” sum.

## Types of Series

### 1. Arithmetic Series

An arithmetic series is the sum of the terms in an arithmetic sequence.

For instance, consider the arithmetic sequence: 4, 5, 7, 9, ... 21.

Let  $S$  be the sum of the terms of this arithmetic sequence. Therefore, we have

$$S = 4 + 5 + 7 + 9 + \dots + 21$$

From here, we can see that  $S$  is an example of an arithmetic series.

Note that in an arithmetic series, there must be a **definite** number of terms. In other words, the addends must be finite. In our previous example which is  $S = 4 + 5 + 7 + 9 + \dots + 21$ , we are just adding the terms from 4 to 21. This allows us to have an “exact” value of  $S$ .

If we don't specify a definite number of terms, we're adding an infinite number of terms.

Suppose that instead of  $S = 4 + 5 + 7 + 9 + \dots + 21$ , we just write it as  $S_2 = 4 + 5 + 7 + 9 + \dots$ .

Since  $S_2$  has no definite end, we're adding an infinite number of terms, and the sum will be larger and larger without bounds.



In short, **an arithmetic series must have a definite number of addends.**

So, to formally define an arithmetic series, we express it mathematically as follows:

$$S_n = a_1 + a_2 + a_3 + \dots + a_n$$

Such that  $a_1, a_2, a_3, \dots, a_n$  are terms of an arithmetic sequence and  $a_n$  represents the end of the sequence.

**Sample Problem:** Which of the following is/are arithmetic serie(s)?

1.  $7 + 11 + 16 + 22 + \dots + 320$
2.  $6 + 12 + 18 + 24 + \dots + 120$
3.  $9 + 8 + 7 + 6 + \dots + (-1)$

**Solution:**

1. This is not an arithmetic series. Note that the addends do not satisfy the condition to become an arithmetic sequence because they have no common difference.
  - $11 - 7 = 4$
  - $16 - 11 = 5$
  - $22 - 16 = 6$
2. This is an arithmetic series since the addends are terms of an arithmetic sequence with a common difference of 6.
3. This is an arithmetic series since the addends are terms of an arithmetic sequence with a common difference of -1.

**Formula for the Sum of an Arithmetic Series**

To find the sum of an arithmetic series, we can use two formulas depending on the given situation.



**Formula #1:  $S_n = n/2[2a_1 + (n - 1)d]$**

**Formula #2:  $S_n = n/2(a_1 + a_n)$**

Such that:

- $a_1$  is the first term
- $n$  is the number of terms being added in the series
- $a_n$  is the  $n$ th term of the sequence. If the sequence has a specified last term, then  $a_n$  is also the last term.
- $d$  is the common difference between the terms or addends in the series

We use the first formula when the sequence has no specified last term. For example, if we want to find the sum of the first 50 terms of  $2 + 4 + 6 + 8 + \dots$ , then we use the first formula. On the other hand, the second formula is used when the last term of the sequence is specified. For example, we can use the second formula to compute the sum of  $1 + 2 + 3 + \dots + 50$ .

It's easy to derive the formulas above. However, we are not going to discuss this anymore because it is beyond the scope of our reviewer.

To further understand how to use these formulas, let us solve some sample questions.

**Sample Problem 1:** What is the sum of all whole numbers from 1 to 100?

**Solution:** By writing the problem as a mathematical sentence, we'll obtain:

$$1 + 2 + 3 + 4 + \dots + 100$$

It's clear here that we are dealing with an arithmetic series since the terms have a common difference of 1. Since the sequence has a specified last term (which is 100), then it's advisable to use the second formula:

- Since the first term of the series is 1, we have  $a_1 = 1$
- The last term of the series is 100, so  $a_n = 100$
- There are 100 whole numbers from 1 to 100, so we have  $n = 100$

Let us insert these values into the formula for the sum of an arithmetic series.

$$\begin{aligned}S_n &= \frac{n}{2} [a_1 + a_n] \\S_n &= \frac{100}{2} [1 + 100] \\S_n &= 50(101) \\S_n &= 5050\end{aligned}$$

Therefore, the sum of numbers from 1 to 100 is 5050. If you have a lot of spare time, you can verify this by doing the addition process manually.

**Sample Problem 2:** What is the sum of  $4 + 6 + 8 + 10 + \dots + 32$ ?

**Solution:** Since the given arithmetic series above has a definite last term (which is 32), then we will use formula #2.

$$S_n = \frac{n}{2} [a_1 + a_n]$$

The first term of the given series is 4. So we have  $a_1 = 4$ . Meanwhile, the last term of the sequence is 32, so we have  $a_n = 32$ .

For the value of  $n$ , we need to determine the number of terms in the series  $4 + 6 + 8 + 10 + \dots + 32$ . We can do this by going back to our formula for the  $n$ th term of an arithmetic sequence. You may manually count the number of terms if you don't want to perform some algebra.

Let us do the algebra stuff by using the formula:

$$a_n = a_1 + (n - 1)d$$

We have  $a_1 = 4$ ,  $a_n = 32$ , and  $d = 2$ . Let us insert these values into the formula above and then solve for the value of  $n$ :

$$a_n = a_1 + (n - 1)d$$

$$32 = 4 + (n - 1)2 \quad \text{By substitution}$$

$$32 = 4 + 2n - 2 \quad \text{Distributive property}$$

$$32 = 2 + 2n \quad \text{Combining like terms}$$

$$-2 + 32 = 2n \quad \text{Transposition method}$$

$$30 = 2n$$

$$30/2 = 2n/2 \quad \text{Dividing both sides by 2}$$

$$n = 15$$

Since we have computed that  $n = 15$ , it means that there are 15 terms from 4 to 32 in the series  $4 + 6 + 8 + 10 + \dots + 32$ .

Our variables for the formula to identify the sum of the arithmetic series are complete.

We have  $a_1 = 4$ ,  $a_n = 32$ ,  $n = 15$ , and  $d = 2$ . Let us substitute these values in the formula:

$$S_n = \frac{n}{2} [a_1 + a_n]$$

$$S_n = \frac{15}{2} [4 + 32]$$

$$S_n = \frac{15}{2} [36]$$

$$S_n = \frac{15}{2} \times \frac{36}{1} = \frac{15 \times 36}{2} = 270$$

Hence, the sum of the arithmetic series  $4 + 6 + 8 + 10 + \dots + 32$  is 270.

**Sample Problem 3:** Bea saved PHP 100 on February 1 in her savings account. On February 2, she saved PHP 105. On February 3, she saved PHP 110, and so on. How much money did Bea save at the end of February (assuming it is not a leap year)?

**Solution:** The manner in which Bea saved money during February follows an arithmetic sequence with a common difference of 5.

- February 1 = 100
- February 2 = 105
- February 3 = 110

To determine the total amount of money that Bea saved for the entire month of February, simply add the daily amount of money saved:

Total amount that Bea saved in February =  $100 + 105 + 110 + \dots$

It is pretty clear that the addition sentence above is an arithmetic series. We shall use the first formula since the last term is not specified.

Since 100 is the first term of the series, we have  $a_1 = 100$

Note that we are looking for the total amount of money Bea saved in February. Since it is not a leap year, Bea was able to save money for 28 days. Therefore, we have  $n = 28$ .

Lastly, the common difference is 5. So we have  $d = 5$ .

Let us input these values into the formula for the sum of an arithmetic series.

$$S_n = \frac{n}{2} [2a_1 + (n - 1)d]$$

$$S_n = \frac{28}{2} [2(100) + (28 - 1)5]$$

$$S_n = \frac{28}{2} [200 + (27)5]$$

$$S_n = 14(200 + 135)$$

$$S_n = 14(335)$$

$$S_n = 4690$$

Therefore, Bea saved a total amount of PHP 4,690 for the entire month of February.

## 2. Geometric Series

A geometric series is the sum of the terms of a geometric sequence.

Consider this geometric sequence: 1, 3, 9, 27, ... Suppose that  $S$  represents the sum of the terms of this sequence such that  $S = 1 + 3 + 9 + 27 + \dots$ . Since  $S$  is the sum of the terms of a geometric sequence, then  $S$  is a geometric series.

Unlike with arithmetic series, the terms in a geometric series can either be finite or infinite. A **finite geometric series** has a definite number of terms. This means that a finite geometric series has a "last" term.

For instance, the geometric series  $2 + 4 + 8 + \dots + 256$  is a finite geometric series because it has a definite number of terms. Note that there is the last term in this series which is 256.

On the other hand, an **infinite geometric series** has an infinite number of terms being added. This means it has no “last” term because it has an endless number of addends.

However, not every infinite geometric series has an “exact” sum. For instance, the infinite geometric series  $2 + 4 + 8 + 16 + \dots$  doesn’t have an exact sum. The sum that we will obtain will just get larger and larger without bounds.

An infinite geometric series will have an “exact” sum if it satisfies a special condition. That is, the absolute value of the common ratio ( $r$ ) must be between 0 to 1:

$$0 < |r| < 1$$

For instance, the geometric series  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$  will have an “exact” sum since its common ratio is  $\frac{1}{2}$  ( $\frac{1}{4} \div \frac{1}{2} = \frac{1}{2}$ ). Note that the absolute value of  $\frac{1}{2}$  is between 0 to 1.

Thus, in general, a finite geometric series can be expressed mathematically as:

$$S_n = a_1 + a_2 + a_3 + a_4 + \dots + a_n$$

where  $a_1, a_2, a_3, \dots, a_n$  are terms of a geometric sequence and  $a_n$  is the “last” addend of the series.

If the absolute value of the common ratio of the terms is between 0 and 1 ( $0 < |r| < 1$ ), then the infinite geometric series can be expressed as:

$$S = a_1 + a_2 + a_3 + a_4 + \dots$$

where  $a_1, a_2, a_3, a_4, \dots$  are terms of a geometric sequence with no definite “last” term.

**Sample Problem 1:** Which of the following is a geometric series?

1.  $3 + (-6) + 12 + \dots + (-96)$
2.  $7 + 9 + 11 + \dots 25$
3.  $2 + 1 + \frac{1}{2} + \dots$

**Solution:** The series in A and C are geometric series with a common ratio of  $-2$  and  $\frac{1}{2}$  respectively. Meanwhile, the series in B is an arithmetic series with a common difference of  $2$ .


**Sample Problem 2:** Which of the following infinite geometric series satisfies the condition to have an exact sum?

1.  $9 + 27 + 81 + \dots$
2.  $-1 + 5 + (-25) + 125 + \dots$
3.  $16 + 4 + 1 + \dots$

**Solution:** A and B have a common ratio of  $3$  and  $-5$  respectively. Note that the respective absolute values of these numbers don't fall within the  $0-1$  range. Hence, they do not satisfy the condition of having an exact sum. On the other hand, the common ratio in C is  $\frac{1}{4}$  which falls within the  $0-1$  range. Therefore, this series will have an exact sum.

#### Formula To Find the Sum of a Finite Geometric Series

Shown below is the formula for the sum of a finite geometric series:

The formula is displayed on a light blue background. It is 
$$S_n = \frac{a_1(1 - r^n)}{(1 - r)}$$
 with a small FilipiKnow logo in the top right corner of the background box.

Such that:

- $a_1$  is the first term (or addend) of the geometric series
- $r$  is the common ratio
- $n$  is the number of terms being added

**Sample Problem 1:** Compute the sum of the first 15 terms of the geometric series  $1 + 2 + 4 + 8 + \dots$



**Solution:**

- The first term of the sequence is 1. So we have  $a_1 = 1$
- The common ratio is 2 ( $2 \div 1 = 2$ ). So we have  $r = 2$
- We are looking for the sum of the first 15 terms. Thus, we have  $n = 15$ .

Let us substitute these values for the formula for the sum of a finite geometric series.

$$S_n = \frac{a_1(1 - r^n)}{1 - r}$$

$$S_n = \frac{1(1 - 2^{15})}{1 - 2}$$

$$S_n = \frac{1(1 - 32768)}{-1}$$

$$S_n = \frac{1(-32767)}{-1} = 32767$$

Based on our calculation above, the sum is 32,767.

**Sample Problem 2:** Compute the sum of the first 6 terms of the sequence  $-3 + 12 + (-48) + \dots$

**Solution:**

- The first term of the sequence is -3. So we have  $a_1 = -3$ .
- The common ratio is -4 ( $12 \div (-3) = -4$ ). So we have  $r = -4$
- We are looking for the sum of the first 6 terms. Thus, we have  $n = 6$ .

Let us insert these values into the formula for the sum of a finite geometric series.

$$S_n = \frac{a_1(1 - r^n)}{1 - r}$$

$$S_n = \frac{-3(1 - (-4)^6)}{1 - (-4)}$$

$$S_n = \frac{-3(1 - (4096))}{5}$$

$$S_n = \frac{-3(-4095)}{5} = 2457$$

Based on our calculation above, the sum is 2,457.

### Formula To Find the Sum of an Infinite Geometric Series

Shown below is the formula for the sum of an infinite geometric series.

$$S_n = \frac{a_1(1 - r^n)}{1 - r}$$

$$S_n = \frac{-3(1 - (-4)^6)}{1 - (-4)}$$

$$S_n = \frac{-3(1 - (4096))}{5}$$

$$S_n = \frac{-3(-4095)}{5} = 2457$$

Such that:

- $a_1$  is the first term (or addend) of the geometric series
- $r$  is the common ratio such that  $0 < |r| < 1$

**Sample Problem:** What is the value of  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$ ?

**Solution:** The first term of the sequence is  $\frac{1}{2}$ . So we have  $a_1 = \frac{1}{2}$ .

Meanwhile, the common ratio is  $\frac{1}{2}$  ( $\frac{1}{4} \div \frac{1}{2} = \frac{1}{2}$ )

By substitution:

$$S_{\infty} = \frac{a_1}{1-r} = \frac{\frac{1}{2}}{1-\frac{1}{2}} = \frac{\frac{1}{2}}{\frac{1}{2}} = 1$$


Thus,  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots = 1$

**BONUS: A Geometric Interpretation on Infinite Geometric Series**


We have learned earlier that using the formula for infinite geometric series, the sum of  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + 1/16 + \dots$  is equal to 1. Although this formula is mathematically sound and valid, I know that this is still too abstract and difficult to grasp.

Using [geometry](#), we can provide a more intuitive explanation of why  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + 1/16 + \dots = 1$ .


Imagine we have a  $\frac{1}{2}$  sheet of colored paper placed on a table.




Now, let us place a  $\frac{1}{4}$  sheet of colored paper near the  $\frac{1}{2}$  sheet of colored paper.




Add a  $\frac{1}{8}$  sheet of colored paper.



Add a  $\frac{1}{16}$  sheet of colored paper. We also perform this step for  $\frac{1}{32}$ ,  $\frac{1}{64}$ ,  $\frac{1}{128}$ , and so on.



Doing this method multiple times allows us to obtain a total amount of paper equivalent to 1 whole sheet of colored paper.



This shows that  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots = 1$ .